

Antimonotonous quadratic forms and partially ordered sets

L.A. Nazarova *, A.V. Roiter *, M.N. Smirnova *

* Institute of Mathematics of National Academy of Sciences of Ukraine,
Tereshchenkivska str., 3, Kiev, Ukraine, ind. 01601
E-mail: roiter@imath.kiev.ua

Let P be a bounded set in n -dimensional space $\mathbb{R}_n$, $f(x_1, \dots, x_n) = f(x)$ ($x \in \mathbb{R}_n$) be a continuous function. By the second Weierstrass theorem ([1], s. 163) $\inf\{f(\overline{P})\}$ ($= \inf_{\overline{P}} f(x)$) is reached. We will call a function f *P-faithful*, if $\inf\{f(\overline{P})\}$ is not reached on $\overline{P} \setminus P$ and $\inf\{f(\overline{P})\} > 0$ (i. e. f is positive on $\overline{P}$). Remark that if $n = 1$, $P = (a, b)$, then it is necessary for P -faithfulness that a function is not monotonous.

Further we will suppose that $P = P_n = \{(x_1, \dots, x_n) \mid 0 < x_i \leq 1, i = \overline{1, n}, x_1 + \dots + x_n = 1\}$. If $n > 1$ then $x_i < 1, i = \overline{1, n}$. Then $\overline{P} = \{(x_1, \dots, x_n) \mid 0 \leq x_i \leq 1, i = \overline{1, n}, x_1 + \dots + x_n = 1\}$. In this case the P -faithfulness f is essentially connected with behavior of function on the hyperplane $H_n = \{(x_1, \dots, x_n) \mid x_1 + \dots + x_n = 0\}$.

For differentiable function f put $C^-(f) = \{h \in H_n \setminus \{0\} \mid \frac{\partial f}{\partial x_i}(h) \leq 0, i = \overline{1, n}\}$, $C^+(f) = \{h \in H_n \setminus \{0\} \mid \frac{\partial f}{\partial x_i}(h) \geq 0, i = \overline{1, n}\}$, $C(f) = C^+(f) \cup C^-(f)$. A function f is *antimonotonous* if $C(f) = \emptyset$. If $n = 1$ then $P_1 = (1)$, $H_1 \setminus \{0\} = \emptyset$ and any function is antimonotonous.

In s. 3 (Proposition 1) we prove that any P -faithful quadratic form is antimonotonous and, therefore, in this case antimonotonousness is a generalization of P -faithfulness.

Example 1. A linear function $f = \sum_{i=1}^n a_i x_i$ is antimonotonous only if all $a_i = 0$, i. e. $f = 0$. The quadratic forms $x_1^2 + x_2^2$, $x_1^2 + x_2^2 + x_1 x_2$ are antimonotonous, but the forms $x_1^2 - x_2^2$, $x_1^2 + x_2^2 + x_3^2 + x_1 x_2 + x_1 x_3$ are not.

The problem of effective criterion of antimonotonousness even for quadratic forms is probably difficult.

In this article we solve this problem for quadratic form f_S , attached to (finite) partially ordered set (poset) $S = \{s_1, \dots, s_n\}$: $f_S(x_1, \dots, x_n) = \sum_{s_i \leq s_j} x_i x_j$ [2] under the additional condition of positive semidefiniteness form f_S (i. e. $f_S(x) \geq 0$). Posets with antimonotonous form generalize P -faithful posets, defined in [3] and studied in [3–7] and (it follows from this work) coincide with them not only for positive definite forms, but for positive semidefinite ones.

A direct construction of a vector contained in $C(f_S)$ permits essentially to simplify the proof of criterion of P -faithfulness [3, 5–7] avoiding consideration of many different cases.

We also take out evident formula for calculating of $\inf\{f_S(\overline{P})\}$ for P -faithful S , on base of this formula we give simple proofs of the criterions of finite representativity [8] (see also [9] and tameness [10], (see also [11]) of partially ordered sets.

1. In s. 1 f is a differentiable function, defined on $\mathbb{R}_n$. We call vectors the elements of $\mathbb{R}_n$.

Let $\mathbb{R}_n^+ = \{x \in \mathbb{R}_n \mid x_i > 0, i = \overline{1, n}\}$, $\overline{\mathbb{R}}_n^+ = \{x = (x_1, \dots, x_n) \in \mathbb{R}_n \mid 0 \leq x_i, i = \overline{1, n}; x \neq 0\}$, $P_n = \overline{P}_n \cap \mathbb{R}_n^+$ ($\mathbb{R}^+ = \mathbb{R}_1^+$).

If f_1, f_2 are defined respectively on $\mathbb{R}_m$ and on $\mathbb{R}_n$ we put $(f_1 \oplus f_2)(x_1, \dots, x_m, x_{m+1}, \dots, x_{m+n}) = f_1(x_1, \dots, x_m) + f_2(x_{m+1}, \dots, x_{m+n})$.

We call a twice differentiable function f *concave*, if the following conditions hold:

- a) $\frac{\partial f}{\partial x_i}(0) = 0, i = \overline{1, n}$,
- b) $\frac{\partial^2 f}{\partial x_i \partial x_j} \geq 0, i, j = \overline{1, n}$,

q -concave, $q \in \mathbb{R}^+$, if a), b) and

- c) $\frac{\partial^2 f}{\partial x_i^2} \geq q, i = \overline{1, n}$ hold.

A quadratic form f_S attached to poset S is, in particular, 2-concave.

Remark 1. By Lagrange theorem b) implies I, c) implies II_q.

I) $\frac{\partial f}{\partial x_i}(x_1, \dots, x_{j-1}, x_j + d, x_{j+1}, \dots, x_n) \geq \frac{\partial f}{\partial x_i}(x_1, \dots, x_n), (i, j \in \overline{1, n})$.

II_q) $\frac{\partial f}{\partial x_i}(x_1, \dots, x_{i-1}, x_i + d, x_{i+1}, \dots, x_n) \geq \frac{\partial f}{\partial x_i}(x_1, \dots, x_n) + qd, (i = \overline{1, n})$;

Put $\widehat{C}^-(f) = \left\{x \in \mathbb{R}_n \setminus \{0\} \mid \sum_{i=1}^n x_i \geq 0, \frac{\partial f}{\partial x_i}(x) \leq 0, i = \overline{1, n}\right\}$; $\widehat{C}^+(f) = \left\{x \in \mathbb{R}_n \setminus \{0\} \mid \sum_{i=1}^n x_i \leq 0, \frac{\partial f}{\partial x_i}(x) \geq 0, i = \overline{1, n}\right\}$.

Lemma 1. If f is a concave function then it is antimonotonous if and only if $\widehat{C}^+(f) \cup \widehat{C}^-(f) = \emptyset$.

Let $x \in \widehat{C}^-(f)$ (the case $x \in \widehat{C}^+(f)$ is analogous), $\sum_{i=1}^d x_i = d \in \mathbb{R}^+$. Then $\{x_1 - d, x_2, \dots, x_n\} \in C(f)$ (according to I) if only $x \neq (d, 0, \dots, 0)$. But at the last case $(d, -d, 0, \dots, 0) \in C(f)$. If $y \in C(f)$ then it is clearly that $y \in \widehat{C}^+(f) \cup \widehat{C}^-(f)$. ■

Lemma 2. If f_1 and f_2 are concave then the function $f_1 \oplus f_2$ is antimonotonous if and only if f_1 and f_2 are antimonotonous.

We shall prove that $C(f_1 \oplus f_2) \neq \emptyset$ if and only if either $C(f_1) \neq \emptyset$ or $C(f_2) \neq \emptyset$. If $(x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}) \in C(f_1 \oplus f_2)$ then either $(x_1, \dots, x_{n_1}) \in \widehat{C}^+(f_1) \cup \widehat{C}^-(f_1)$ or $(y_1, \dots, y_{n_2}) \in \widehat{C}^+(f_2) \cup \widehat{C}^-(f_2)$ and by Lemma 1 in the first case $C(f_1) \neq \emptyset$ and in the second case $C(f_2) \neq \emptyset$.

If $x = (x_1, \dots, x_{n_1}) \in C(f_1)$ then $(x_1, \dots, x_{n_1}, \underbrace{0, \dots, 0}_{n_2}) \in C(f_1 \oplus f_2)$ using a); if $(y_1, \dots, y_{n_2}) \in C(f_2)$ then $(\underbrace{0, \dots, 0}_{n_1}, y_1, \dots, y_{n_2}) \in C(f_1 \oplus f_2)$. ■

We say that a nonzero vector $d \in \mathbb{Z}_n$ is *m-Dynkin*) ($1 \leq m \leq n$) for q -concave function f , if 1) $0 \leq \frac{\partial f}{\partial x_m}(d) \leq q$; 2) $\frac{\partial f}{\partial x_j}(d) = 0$ if $j \neq m, j = \overline{1, n}$.

We call a function f *m-isolated*, if $\frac{\partial f}{\partial x_k}(s_m) = 0$ for $1 \leq k \leq n, k \neq m, s_m = (\underbrace{0, \dots, 0}_{m-1}, 1, 0, \dots, 0)$.

Lemma 3. Let f be q -concave not m -isolated function. If there exists m -Dynkin vector then $C(f) \neq \emptyset$.

Let $\sum_{i=1}^n d_i = \bar{d}$. If $\bar{d} \leq 0$ then $d \in \widehat{C}^+(f)$ and $C(f) \neq \emptyset$ by Lemma 1. Let $\bar{d} > 0$. Put $u_j = d_j$ if $j \neq m$ and $u_m = d_m - \bar{d}$, and we prove that $u = (u_1, \dots, u_n) \in C(f)$. It is clear that $u \in H_n$. $\frac{\partial f}{\partial x_j}(u) \leq 0$ if $j \neq m$ according to I and 2, and $\frac{\partial f}{\partial x_m}(u) \leq 0$ according to II_q and 1).

Now we shall prove that $u \neq 0$. If $u = 0$ then $d = \lambda s_m$, $\lambda \neq 0$ (since $d \neq 0$). But then non m -isolateness f implies that $\frac{\partial f}{\partial x_k}(d) \neq 0$, $k \neq m$. ■

Example 2. Let $S = \{s_1, s_2, s_3, s_4, s_5 \mid s_1 < s_i, i = \overline{2, 5}\}$, $f_S = \sum_{i=1}^5 x_i^2 + x_1 \sum_{j=2}^5 x_j$, $d = (-2, 1, 1, 1, 1)$ — i -Dynkin vector for f_S ($i = \overline{1, 5}$). Vectors $(-2, 1, 1, 1, -1)$, $(-2, 1, 1, -1, 1)$, $(-2, 1, -1, 1, 1)$, $(-2, -1, 1, 1, 1)$ and $(-4, 1, 1, 1, 1)$ belong to $C(f_S)$.

Consider P -faithfulness. Put $\text{St}(f) = \{a \in \mathbb{R}_n^+ \mid \frac{\partial f}{\partial x_i}(a) = \frac{\partial f}{\partial x_j}(a), i, j = \overline{1, n}\}$; $\text{St}^+(f) = \{a \in \text{St} \mid \frac{\partial f}{\partial x_i}(a) > 0\}$.

Vector $u \in P_n$ is P -faithful for function f , if $f(u) > 0$ and $w \in \overline{P}_n$ implies $f(u) \leq f(w)$, moreover if $w \notin P_n$, then $f(u) < f(w)$.

Denote $\widetilde{\text{St}}(f)$ the set of P -faithful vectors for f . P -faithfulness of f is equivalent to $\widetilde{\text{St}}(f) \neq \emptyset$.

Lemma 4. $\widetilde{\text{St}}(f) \subseteq \text{St}(f)$ for any f .

Let $n > 1$. We express $x_n = 1 - \sum_{i=1}^{n-1} x_i$, and obtain the function $\widehat{f}(x_1, \dots, x_{n-1}) = f(x_1, \dots, x_{n-1}, 1 - \sum_{i=1}^{n-1} x_i)$. If $u = (u_1, \dots, u_n)$ is P -faithful vector for f then $\widehat{f}(\widehat{u})$ is minimum when $\widehat{u} = (u_1, \dots, u_{n-1})$. $\frac{\partial \widehat{f}}{\partial x_i} = \frac{\partial f}{\partial x_i} + \frac{\partial f}{\partial x_n} \cdot \frac{\partial x_n}{\partial x_i}$; $x_n = 1 - \sum_{i=1}^{n-1} x_i$; $\frac{\partial x_n}{\partial x_i} = -1$ ($i = \overline{1, n-1}$). Therefore $\frac{\partial \widehat{f}}{\partial x_i} = \frac{\partial f}{\partial x_i} - \frac{\partial f}{\partial x_n} = 0$. ■

Let f be a homogeneous function of degree k (i. e. $f(\lambda x_1, \dots, \lambda x_n) = \lambda^k f(x_1, \dots, x_n)$). For homogeneous f , if $k \neq 1$ and $\inf\{f(P)\} > 0$ put $P(f) = \inf\{f(P)\}^{\frac{1}{1-k}}$. In particular, for $k = 2$, $P(f) = \inf\{f(\overline{P})\}^{-1}$.

Lemma 5. Let $f_1(x_1, \dots, x_{n_1})$, $f_2(x_{n_1+1}, \dots, x_{n_2})$ be two homogeneous functions of degree k , $n_1 + n_2 = n$, $\inf\{f_j(P_{n_j})\} > 0$, $j = 1, 2$. Then $P(f_1 \oplus f_2) = P(f_1) + P(f_2)$.

Values of homogeneous function f on $\mathbb{R}_n^+$ are defined by its values on $\overline{P}_n$, namely for $y \in \mathbb{R}_n^+$, $f(y) = \lambda^k f(u)$ where $u \in \overline{P}_n$, $\lambda = \sum_{i=1}^n y_i$, $u = \lambda^{-1}y$. Therefore $\inf\{(f_1 \oplus f_2)(\overline{P}_n)\} = \inf_{0 \leq \lambda \leq 1} [\lambda^k \inf\{f_1(\overline{P}_{n_1})\} + (1 - \lambda)^k \inf\{f_2(\overline{P}_{n_2})\}]$. Put $\inf\{f_1(\overline{P}_{n_1})\} = a$, $\inf\{f_2(\overline{P}_{n_2})\} = b$.

Consider function $\Phi_{ab}(\lambda) = a\lambda^k + b(1 - \lambda)^k$, $a > 0$, $b > 0$ and find $\inf_{0 \leq \lambda \leq 1} \Phi_{ab}(\lambda)$. The derivative of $\Phi_{ab}(\lambda)$ with respect to λ (consider u and v to be constants) is $(\Phi_{ab}(\lambda))'_\lambda = ka\lambda^{k-1} - kb(1 - \lambda)^{k-1}$. Let $\bar{\lambda}$ be a positive root of equation $(\Phi_{ab}(\lambda))'_\lambda = 0$. Substitute $\bar{\lambda}$ and obtain $a\bar{\lambda}^{k-1} = b(1 - \bar{\lambda})^{k-1}$. Raise the both parts of this equality to $\frac{1}{k-1}$ power and obtain

$a^{\frac{1}{k-1}}\bar{\lambda} = b^{\frac{1}{k-1}}(1 - \bar{\lambda})$, $\bar{\lambda} = \frac{b^{\frac{1}{k-1}}}{a^{\frac{1}{k-1}} + b^{\frac{1}{k-1}}}$. Thus, $\inf_{0 \leq \lambda \leq 1} \Phi_{ab}(\lambda) = \min\{\Phi_{ab}(0), \Phi_{ab}(1), \Phi_{ab}(\bar{\lambda})\} = \min\{a, b, \Phi_{ab}(\bar{\lambda})\}$.

We will prove that $\Phi_{ab}(\bar{\lambda}) < \Phi_{ab}(0) = b$. Really, $a\bar{\lambda}^{k-1} = b(1 - \bar{\lambda})^{k-1}$. $\Phi_{ab}(\bar{\lambda}) = a\bar{\lambda}^k + b(1 - \bar{\lambda})^k = b(1 - \bar{\lambda})^{k-1}\bar{\lambda} + b(1 - \bar{\lambda})^k = b(1 - \bar{\lambda})^{k-1}$. $\Phi_{ab}(\bar{\lambda}) < b$, since $a > 0$, $b > 0$, $0 < \bar{\lambda} < 1$. Analogously $\Phi_{ab}(\bar{\lambda}) = a\bar{\lambda}^{k-1} < \Phi_{ab}(1) = a$. Therefore $\inf\{(f_1 \oplus f_2)(\bar{P}_n)\} = \Phi_{ab}(\bar{\lambda}) = \frac{ab}{\left(a^{\frac{1}{k-1}} + b^{\frac{1}{k-1}}\right)^{k-1}}$.

Let us return to $P(f_1 \oplus f_2)$, $P(f_1)$, $P(f_2)$. We have $P(f_1) = a^{\frac{1}{1-k}}$, $P(f_2) = b^{\frac{1}{1-k}}$, $P(f_1 \oplus f_2) = \left(\frac{ab}{\left(a^{\frac{1}{k-1}} + b^{\frac{1}{k-1}}\right)^{k-1}}\right)^{\frac{1}{1-k}} = b^{\frac{1}{1-k}} + a^{\frac{1}{1-k}}$. ■

Corollary 1. *Under the conditions of the lemma $f_1 \oplus f_2$ is P -faithful if and only if f_1 and f_2 are P -faithful.*

2. Further $f = \sum_{i,j=1}^n a_{ij}x_i x_j$ ($a_{ij} = a_{ji}$) is a quadratic form over field $\mathbb{R}$; $A = (a_{ij})$ is a symmetric matrix of quadratic form f . $\frac{\partial^2 f}{\partial x_i \partial x_j} = 2a_{ij}$. f is a 2-concave if $a_{ii} \in \mathbb{N}$, $a_{ij} + a_{ji} \in \mathbb{N}_0$ ($i, j = \overline{1, n}$). For $x = (x_1, \dots, x_n) \in \mathbb{R}_n$ (for fixed f) put

$$x'_i = \frac{\partial f}{\partial x_i}(x_1, \dots, x_n) = 2 \sum_{j=1}^n a_{ji}x_j, \quad x' = (x'_1, \dots, x'_n) = 2xA.$$

We will use the following identity, which is easy to check:

$$f(u + v) = f(u) + f(v) + \sum_{i=1}^n u'_i v_i, \quad u, v \in \mathbb{R}_n, \quad \text{that implies} \quad (1)$$

$$\sum_{i=1}^n u'_i v_i = \sum_{i=1}^n v'_i u_i, \quad f(u + \varepsilon v) = f(u) + \varepsilon^2 f(v) + \varepsilon \sum_{i=1}^n u_i v'_i, \quad \varepsilon \in \mathbb{R}. \quad (2)$$

Put in (1) $u = v$, and obtain (see. [1], s. 178):

$$f(u) = \frac{1}{2} \sum_{i=1}^n u_i u'_i. \quad (3)$$

By means of (3) we can reformulate Lemma 4 by the following way.

Lemma 4'. $\tilde{\text{St}}(f) \subset \text{St}^+(f)$ for quadratic form f .

Put $\tilde{C}(f) = \{(v_1, \dots, v_n) \in C(f) \mid (v'_1, \dots, v'_n) \neq 0\}$. Since $\frac{\partial f}{\partial x_i}(-x) = -\frac{\partial f}{\partial x_i}(x)$ then if $C(f) \neq \emptyset$, then $C^-(f) \neq \emptyset$ and $C^+(f) \neq \emptyset$. Therefore, choosing a vector $v \in C(f) \neq \emptyset$, further we will suppose that $v \in C^-(f)$.

Proposition 1. *For any quadratic form f 1) at least one of the sets $\text{St}(f)$ and $\tilde{C}(f)$ is empty 2) at least one of the sets $C(f)$ and $\tilde{\text{St}}(f)$ is empty.*

1) Let $u \in \text{St}(f)$, $v \in \tilde{C}(f)$. Then $\sum_{i=1}^n u'_i v_i = u'_1 \sum_{i=1}^n v_i = 0$. In the other hand, if $v'_j < 0$ then $\sum_{i=1}^n u_i v'_i < 0$ (since $v'_i \leq 0$, $u_i > 0$, $i = \overline{1, n}$), that contradicts (2).

2) Let $u \in \tilde{\text{St}}(f)$, $v \in C(f)$. If $v \in \tilde{C}(f)$ then 1) implies the statement. Let $v \in C(f) \setminus \tilde{C}(f)$, i. e. $v'_i = 0$ for $i = \overline{1, n}$. Then $f(v) = 0$ ((3)) and so $f(u + \varepsilon v) = f(u)$ for any ε . Put $|\varepsilon| = \min_i \frac{u_i}{|v_i|}$. The sign of ε is opposite sign of one of those v_i , for which the minimum is reached. Then $u + \varepsilon v \in \overline{P}_n \setminus P_n$, that contradicts P -faithfulness of u . ■

Corollary 2. *P -faithful quadratic form is antimonotonous.*

Example 3. Let $f = \sum_{i=1}^4 x_i^2 + (x_1 + x_2)(x_3 + x_4)$. Then

$$A = \begin{pmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 1 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 1 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 & 1 \end{pmatrix}, \quad \text{St}(f) \ni (1, 1, 1, 1), \quad C(f) \ni (1, 1, -1, -1).$$

Proposition 1 implies $\tilde{\text{St}}(f) = \emptyset$, $\tilde{C}(f) = \emptyset$.

In this example $|A| = 0$.

Proposition 2. *If $|A| \neq 0$ then one of the sets $C(f)$, $\text{St}(f)$ is not empty, but the other one is empty.*

At first suppose that $\emptyset \neq C(f) \ni v$ and $\emptyset \neq \text{St}(f) \ni u$. If $v \in \tilde{C}(f)$ then $\text{St}(f) = \emptyset$ by Proposition 1. If $v \in C(f) \setminus \tilde{C}(f)$ then $v' = 0$, and $vA = \frac{1}{2}v' = 0$. Hereof $v = 0$ that contradicts $v \in H_n \setminus \{0\}$ (the definition of $C(f)$).

Now we prove that either $\text{St}(f) \neq \emptyset$, or $C(f) \neq \emptyset$. Let $e_n = (1, \dots, 1) \in \mathbb{R}_n$, $y = e_n A^{-1}$, $yA = e_n$. If $y \in \mathbb{R}_n^+$ or $-y \in \mathbb{R}_n^+$, then $y \in \text{St}(f)$. If $\{y, -y\} \cap \mathbb{R}_n^+ = \emptyset$ then either for certain k $y_k = 0$, or for certain s and t $y_s < 0$, $y_t > 0$. It is easy to see that in both cases there exists $w \in \overline{\mathbb{R}}_n^+$ such that $wy^T \left(= \sum_{i=1}^n w_i y_i \right) = 0$ (at the first case we can put $w_k > 0$, $w_i = 0$ for $i \neq k$, at the second case $w_s = y_t$, $w_t = -y_s$, $w_i = 0$ for $i \notin \{s, t\}$, $i = \overline{1, n}$). We prove that $v = -wA^{-1} \in C(f)$. $-v' = wA^{-1}A = w \in \overline{\mathbb{R}}_n^+$, and so $v'_i \leq 0$. $v \neq 0$, since $w \neq 0$ and $|A| \neq 0$. It remains to prove that $v \in H_n$, that is equivalent to $ve_n^T = 0$. $ve_n^T = -wA^{-1}e_n^T$; $y^T = (A^{-1})^T e_n^T = A^{-1}e_n^T$ (since $A^T = A$). Therefore $-wA^{-1}e_n^T = -wy^T = 0$. ■

Proposition 3 (see [7], p. II, remark to theorem 1). 1) *If $\tilde{\text{St}}(f) \neq \emptyset$, then f is positive definite.* 2) *If f is positive definite then $\tilde{\text{St}}(f) = \text{St}(f) \cap P_n$ (and therefore $\tilde{\text{St}}(f) = \emptyset$ if and only if $\text{St}(f) = \emptyset$).*

1) We suppose contrary: $f(v) \leq 0$ ($v \neq 0$), $u \in \tilde{\text{St}}(f)$. a) At first we suppose that $v \in H_n$, i. e. $\sum_{i=1}^n v_i = 0$ and $f(v) < 0$. Then $f(u + \varepsilon v) = f(u) + \varepsilon^2 f(v) + \varepsilon \sum_{i=1}^n u'_i v_i$. By Lemma 4 $u \in \text{St}(f)$ and, therefore, $\varepsilon \sum_{i=1}^n u'_i v_i = 0$, i. e. $f(u + \varepsilon v) = f(u) + \varepsilon^2 f(v)$. Since $f(v) < 0$ then $f(u + \varepsilon v) < f(u)$, that contradicts P -faithfulness of u .

b) Now let $v \in H_n$, $f(v) = 0$. Then $f(u + \varepsilon v) = f(u)$ for any ε . Put $\varepsilon = \min_i \frac{u_i}{|v_i|}$. The sign of ε is opposite to the sign of one of those v_i , for which this minimum is reached. Then $u + \varepsilon v \in \overline{P}_n \setminus P_n$, again we have the contradiction with P -faithfulness of u .

c) Let finally $\sum_{i=1}^n v_i \neq 0$. We can admit $\sum_{i=1}^n v_i = 1$. Put $w = u - v$. (2) for $\varepsilon = -1$ and Lemma 4 imply $f(w) = f(u) + f(v) - u'$, $u' = u'_i$, $i = \overline{1, n}$. (3) implies $f(u) = \frac{u'_1}{2}$ and, therefore, $f(w) = f(v) - \frac{u'_1}{2}$. By Lemma 4' $u' > 0$, $f(w) < 0$; $w \in H_n$. Hereat $w \neq 0$, since if $w = 0$ then $u = v$, according to (3), but $f(v) \leq 0$. Thus, we reduced c) to a).

2) Let $u \in \text{St}(f) \cap P_n$, $v \in \overline{P}_n$, $v \neq u$. $u \neq 0$ since $u \in \text{St}(f)$ so $f(u) > 0$. We will prove that $f(u) < f(v)$. $0 < f(u - v) \stackrel{\text{formula (2)}}{=} f(u) + f(v) - \sum_{i=1}^n u'_i v_i \stackrel{\text{formula (3)}}{=} \frac{u'_1}{2} + f(v) - u' = f(v) - \frac{u'_1}{2} = f(v) - f(u)$, i. e. $f(v) > f(u)$. ■

3. Further we will consider 2-concave form f_S for poset $S = \{s_1, \dots, s_n\}$, $f_S = \sum_{s_i \leq s_j} x_i x_j$. Put $C(S) = C(f_S)$. S is *antimonotonous* if f_S is antimonotonous.

A poset S is *P-faithful*, if $\tilde{\text{St}}(f_S) \neq \emptyset$. (It is equivalent to the definition of P -faithfulness of the poset from [3]) In this case Proposition 1 implies $C(S) = \emptyset$. Remark that $\inf\{f_S(\overline{P})\} > 0$ since $a_{ij} \geq 0$, $i, j = \overline{1, n}$, $A \neq (0)$.

Hasse quiver (orgraph) $Q(S)$ of poset S is a quiver, whose vertices are elements of S and two vertices are connected by an arrow $s_i \rightarrow s_j$ if $s_i < s_j$ and there is no $s_k \in S$ such that $s_i < s_k < s_j$. Drawing lines (edges) instead of arrows, we obtain (nonoriented) graph Hasse $\Gamma(S)$ of partially ordered set S . Finite poset S is usually depicted by a diagram, i. e. by graph $\Gamma(S)$ assuming that lesser element is drawn below than greater.

We denote elements of poset S and corresponding elements of $Q(S)$ and $\Gamma(S)$ by the same symbol.

Put $\text{St}(S) = \text{St}(f_S)$, $\tilde{\text{St}}(S) = \tilde{\text{St}}(f_S)$. Let $\mathcal{T}_S(x_1, \dots, x_n) = \sum_{i=1}^n x_i^2 - \sum_{s_i \rightarrow s_j} x_i x_j$ be a quadratic Tits form of graph $\Gamma(S)$ (the second sum is taken by all edges of graph $\Gamma(S)$). We denote matrix of the form $\mathcal{T}_S$ either $\mathcal{A}$ or $\mathcal{A}(S)$.

Let Q be a quiver without loops and parallel (i. e. having the same origin and the same terminus) pathes with vertices $s_1, \dots, s_n$. $\tilde{Q}$ is a matrix, in which $\tilde{Q}_{ij} = 1$, if there is an arrow from s_i to s_j , $\tilde{Q}_{ij} = 0$ in the opposite case ($i, j = \overline{1, n}$). Then $(\tilde{Q}^t)_{ij}$ is equal to the number of pathes of length t from s_i to s_j , $\tilde{Q}^n = 0$. If, moreover, $Q = Q(S)$ then $A = \frac{1}{2}[(E + \tilde{Q} + \dots + \tilde{Q}^{n-1}) + (E + \tilde{Q} + \dots + \tilde{Q}^{n-1})^T]$ (A is the matrix of f_S). It is easy to see that $(E + \tilde{Q} + \dots + \tilde{Q}^{n-1}) = (E - \tilde{Q})^{-1}$. Put $E - \tilde{Q} = \hat{Q}$, $|\hat{Q}| = 1$. $\mathcal{A} = \frac{1}{2}(\hat{Q} + \hat{Q}^T)$. $A = \frac{1}{2}(\hat{Q}^{-1} + (\hat{Q}^{-1})^T)$.

Proposition 4 ([4])¹ *If there are no parallel pathes in $Q(S)$ then the forms $\mathcal{T}_S$ and f_S are equivalent over $\mathbb{Z}$.*

Really, $\hat{Q}^{-1} \mathcal{A}_T (\hat{Q}^{-1})^T = \frac{1}{2} \hat{Q}^{-1} (\hat{Q} + \hat{Q}^T) (\hat{Q}^{-1})^T = \frac{1}{2} [(\hat{Q}^{-1})^T + \hat{Q}^{-1}] = A$. ■

Propositions 1, 2, 3, 4 imply

¹A.I. Sapelkin in [6] called this statement as Zeldich lemma. M.V. Zeldich in [7] called it “important and surprising”, with what authors quite agree, in spite of brevity of the proof


Corollary 3. *Let $\Gamma(S)$ be acyclical and at least one of the forms f_S , $\mathcal{T}_S$ be positive definite (it holds if $\Gamma(S)$ is a Dynkin graph, see s. 4). Then the other form is positive definite as well, and the following statements are equivalent:*

- a) S is antimonotonous;
- b) S is faithful;
- c) $\text{St}(S) \neq \emptyset$.

We denote $I(s_i)$ for $s_i \in S$ the number of edges of graph $\Gamma(S)$, having s_i its terminus.

s_i is a *terminal point* if $I(s_i) \leq 1$, s_i is a *branch point* if $I(s_i) \geq 3$. s_i is a *junction point*, if it is either a terminus of at least two arrows, or an origin of at least two arrows of quiver $Q(S)$. We denote $S^\times$ the set of junction points.

Example 4.



Here $S^\times = S$, $C(S) \cup \tilde{\text{St}}(S) = \emptyset$. Really, values of f_S can be negative ($f_S(1, 1, 1, -1, -1, -1) = -2$), consequently, by Proposition 3, $\tilde{\text{St}}(S) = \emptyset$, but $\text{St}(S) \ni (1, 2, 1, 1, 2, 1)$, $|A| = -48$. Proposition 2 implies $C(S) = \emptyset$.

We will assume further (except Appendix) graph $\Gamma(S)$ to be connected.

We remind that a cycle in graph Γ is a sequence $\{s_1, \dots, s_m\}$ of different vertices s_i of graph Γ where $m \geq 3$ such that a vertex s_i is connected with s_{i+1} for $i = \overline{1, m}$, and a vertex s_m is connected with s_1 . We call a cycle $\{s_1, \dots, s_m\}$ *simple*, if there are no other vertices between $s_1, \dots, s_m$. We call graph Γ and poset S such that $\Gamma(S) = \Gamma$ *cyclical*, if Γ contains a cycle, and *acyclical* in opposite case. It is easy to see that a cyclical graph contains a simple cycle, and, correspondingly, a cyclical poset S contains a subset S' such that $\Gamma(S') = A_m$. It is clear that if S is acyclical then $Q(S)$ has no parallel pathes.

If $\Gamma(S) = \Gamma(\vec{S})$ then $Q(\vec{S})$ can be obtained from $Q(S)$ by “reorientation” (i. e. by alternation of direction) of several arrows. If $\Gamma(S)$ is acyclic and quiver $\vec{Q}$ is obtained by reorientation of arrows from $Q(S)$, then there exists $\vec{S}$ such that $\vec{Q} = Q(\vec{S})$.

We call quiver $Q = Q(S)$ and poset S *standard*, if $I(s_i) = 2$ implies that s_i is origin of one arrow and terminus also of one arrow, and $I(s_i) \neq 2$ implies that s_i is either origin of $I(s_i)$ arrows or terminus of $I(s_i)$ arrows ($i = \overline{1, n}$). It is easy to see that exactly one standard poset is attached to each acyclical graph up to antiisomorphism. If S^* is antiisomorphic to S then $\Gamma(S) = \Gamma(S^*)$, and $Q(S^*)$ is obtained from $Q(S)$ by reorientation of all arrows.

If φ is an arrow of $Q(S)$, then we denote $S(\varphi)$ poset, obtaining from S by overturn of an arrow φ . A_φ is matrix of $f_{S(\varphi)}$. It is clear that $\mathcal{A}(S(\varphi)) = \mathcal{A}(S)$.

We call a point $s_m \in S$ *Dynkin*, if there exists m -Dynkin vector for form f_S .

Remark 2. Function f_S is m -isolated in sense of s.1, if s_m is not comparable with other points of S . Therefore for connected S the condition of non-isolateness of f_S in Lemma 3 holds automatically.

Lemma 6. *Let $\Gamma(S)$ be acyclical, $s_i \xrightarrow{\varphi} s_j \in Q(S)$; $d \neq 0$ is such a vector that $d'_i = 2(dA)_i = 0$, $d'_j = 2(dA)_j = 0$. Then there exists a vector $\hat{d} \neq 0$ such that $dA = \hat{d}A_\varphi$.*

It follows from the proof of Proposition 4 that $\tilde{Q}^{-1}\mathcal{A}(\tilde{Q}^{-1})^T = A$, $\tilde{Q}_\varphi^{-1}\mathcal{A}_\varphi(\tilde{Q}_\varphi^{-1})^T = A_\varphi$, $\tilde{Q}_\varphi^{-1}\tilde{Q}A\tilde{Q}^T(\tilde{Q}_\varphi^T)^{-1} = A_\varphi$ ($\mathcal{A}_\varphi = \mathcal{A}$). Put $\hat{d} = d\tilde{Q}^{-1}\tilde{Q}_\varphi$, ($\hat{d} \neq 0$), $\hat{d}A_\varphi = d\tilde{Q}^{-1}\tilde{Q}_\varphi\tilde{Q}_\varphi^{-1}\tilde{Q}A\tilde{Q}^T(Q_\varphi^T)^{-1} = dA\tilde{Q}^T(\tilde{Q}_\varphi^T)^{-1}$.

Put (in view of $d'_i = d'_j = 0$), $dA = \sum_{k \notin \{i,j\}} \alpha_k s_k = b$. We shall prove that $b\tilde{Q}^T(\tilde{Q}_\varphi^T)^{-1} = b$, i. e. that $s_k\tilde{Q}^T(\tilde{Q}_\varphi^T)^{-1} = s_k$, i. e. that $s_k\tilde{Q}^T = s_k\tilde{Q}_\varphi^T$, that follows from the definition of $\tilde{Q}$ and from $k \notin \{i,j\}$. ■

Lemma 7. *If $\Gamma(S)$ is acyclical, s_t is a Dynkin terminal point of S then it is Dynkin point for poset $\vec{S}$ if $\Gamma(\vec{S}) = \Gamma(S)$.*

If $\vec{S} = S(\varphi)$ then the statement follows from Lemma 6. Considering the general case ($\vec{S}$ is not $S(\varphi)$), we remark firstly that if S^* and S are antiisomorphic then $f_S = f_{S^*}$ and s_t is a Dynkin point also for S^* .

We denote ψ (resp. $\hat{\psi}$) unique arrow $Q(S)$ (resp. $Q(\vec{S})$) for which s_t is either terminus or origin. Then the condition $s_t \notin \{s_i, s_j\}$ is equal to $\varphi \neq \psi$.

Without loss of generality suppose that ψ in $Q(S)$ and $\hat{\psi}$ in $Q(\vec{S})$ have the same orientation, otherwise pass to $\vec{S}^*$. In this case we can pass from S to $\vec{S}$, returning several arrows, different from ψ , and, therefore, partial cases $\vec{S} = S(\varphi)$ (Lemma 6) and $\vec{S} = S^*$ considered by us imply the statement of lemma. ■

4. Let Γ be a connected acyclical graph with one branch point and three terminal points. Γ is a union of three chains A_{n_1} , A_{n_2} , A_{n_3} , intersecting in a branch point s_1 . $\Gamma = A_{n_1} \cup A_{n_2} \cup A_{n_3}$, $A_{n_1} \cap A_{n_2} = A_{n_1} \cap A_{n_3} = A_{n_2} \cap A_{n_3} = \{s_1\}$, $|A_{n_j}| = n_j$, $j = \overline{1,3}$, $|\Gamma| = n_1 + n_2 + n_3 - 2$. We will denote Γ by $\Gamma(n_1, n_2, n_3)$ (graph does not change if we permute n_j).

All Dynkin graphs besides A_n (i. e. D_n , E_6 , E_7 , E_8) and extended Dynkin graphs $\tilde{E}_6$, $\tilde{E}_7$, $\tilde{E}_8$ have the form $\Gamma(n_1, n_2, n_3)$. It is well-known that $\Gamma(n_1, n_2, n_3)$ is a Dynkin graph if and only if $n_1^{-1} + n_2^{-1} + n_3^{-1} > 1$, and is an extended Dynkin graph when $n_1^{-1} + n_2^{-1} + n_3^{-1} = 1$.

Namely, $\Gamma(n_1, n_2, n_3)$ is E_6 , E_7 , E_8 , D_n , if respectively $(n_1, n_2, n_3) = (3, 3, 2)$; $(2, 4, 3)$; $(2, 3, 5)$; $(1, 1, n-2)$. $\Gamma(m_1, m_2, m_3)$ is respectively $\tilde{E}_6$, $\tilde{E}_7$, $\tilde{E}_8$. for $(m_1, m_2, m_3) = (3, 3, 3)$; $(2, 4, 4)$; $(2, 3, 6)$.

We fixed here the numeration of m_j and n_j so that $m_1 \leq m_2 \leq m_3$ and for E_n ($n = 6, 7, 8$), $n_1 = m_1$, $n_2 = m_2$, $n_3 = m_3 - 1$.

Remark, that (in all cases) m_3 is divided by m_1 and m_2 .

Proposition 5. *If $\Gamma(S) = \Gamma(n_1, n_2, n_3)$ is Dynkin graph or extended Dynkin graph (i. e. D_n , E_6 , E_7 , E_8 , $\tilde{E}_6$, $\tilde{E}_7$ or $\tilde{E}_8$) then S contains terminal Dynkin point.*


By Lemma 7 we can, without loss of generality suppose S to be standard.

For any $\Gamma(m_1, m_2, m_3)$ ($I(s_1) = 3$) we construct a vector $\tilde{d}$, putting $\tilde{d}_1 = -m_3$; $\tilde{d}_i = \frac{m_3}{m_j}$, for $s_i \in A_{n_j}$, $i \neq 1$. It is easy to see that $\tilde{d}'_i = 0$, for $i \neq 1$, $\tilde{d}'_1 = m_3(1 - m_1^{-1} - m_2^{-1} - m_3^{-1})$. If, moreover, $\Gamma(S)$ is extended Dynkin graph then $\tilde{d} \in \mathbb{Z}_n$ and $\tilde{d}'_1 = 0$, i. e. $\tilde{d}$ is in this case a i -Dynkin vector for any i .

Let $\Gamma(S)$ be E_n , $|S| = n$. $\vec{S}$ is such standard poset that $\Gamma(\vec{S}) = \tilde{E}_n$, $|\vec{S}| = n+1$, $S \subset \vec{S}$, $\vec{S} \setminus S = \{s_{n+1}\} \subset A_{n_3}$. We construct for S a Dynkin vector d , modifying Dynkin vector $\tilde{d}$


for $\Gamma(\tilde{S})$. Put $d_i = \tilde{d}_i$ for $i < n$, and $d_n = 2$ ($= \tilde{d}_n + \tilde{d}_{n+1}$), $d'_n = 1$; ($d'_i = 0$, $i = \overline{1, n-1}$).

Let




then $w = (w_1, \dots, w_n)$ where $w_1 = -2$, $w_2 = w_3 = 1$, $w_n = 2$, $w_i = 0$, $i \notin \{1, 2, 3, n\}$ is n -Dynkin vector s_n ($w'_n = 2$). ■

We write out evidently Dynkin vector for standard S if $\Gamma(S) = E_6, E_7, E_8$




Example 5. Dynkin vector for standard poset S such that $\Gamma(S) = \tilde{D}_n$, $n > 4$ (for $n = 4$ see Example 2) has the following form (all points are Dynkin points)



5. Let poset $V = h_1 \blacktriangleleft h^+ \blacktriangleright h_2$, poset $W^{2k} = \{s_1^-, \dots, s_k^-, s_1^+, \dots, s_k^+ \mid s_i^- < s_i^+, s_i^- < s_{i+1}^-, s_k^- < s_1^+, s_k^+ < s_1^-\}$

$s_{i+1}^-, s_k^- < s_1^+, i = \overline{1, k}\}$, $k > 1$, in particular, $W^4 =$



(see Example 3).

Lemma 8. If poset S is cyclical and each $S' \subset S$ is acyclical then S is either V , or W^{2k} ($k \geq 2$).

Without loss of generality suppose that $\Gamma(S)$ is a simple cycle $\tilde{A}_n =$



Let $S \setminus S^\times \ni s$, then $s^- < s < s^+$ and in $\Gamma(S)$ s^- and s^+ are connected with s by edges. Then consider $S' = S \setminus \{s\}$. If there is an edge $s^- - s^+$ in graph $\Gamma(S')$ then $\Gamma(S')$ is a cycle (that contradicts the condition of lemma).

If s^- and s^+ are not connected by edge in $\Gamma(S')$ then there is a point $\bar{s} \neq s$ in S such that $s^- < \bar{s} < s^+$, $\bar{s} \not\propto s$ (i. e. $\bar{s}$ and s are not comparable) because in the opposite case s^- and s^+ should be not connected with s in $\Gamma(S)$. And so $\{s^-, s^+, s, \bar{s}\} = V$ and $S = V$.

So, if $S \neq V$ then $S^\times = S$ and then (since $\Gamma(S) = \tilde{A}_n$) it is easy to see that $S = W^{2k}$ for some $k > 1$. ■

Lemma 9. *If $S \supseteq V$ and $S \not\supseteq W^4$ then $C(S) \neq \emptyset$.*

An arbitrary vector $v \in \mathbb{R}_n$ can be considered as a function on S with values in $\mathbb{R}$.

Let $v : S \rightarrow \mathbb{R}$. $v(h^-) = v(h^+) = -1$, $v(h_1) = v(h_2) = 1$, $v(t) = 0$ for $t \in S \setminus V$; $v \in H_n$. We prove that $v'(s) \leq 0$ for $s \in S$. $v'(h^-) = v'(h^+) = -1$, $v'(h_1) = v'(h_2) = 0$. If t is not comparable neither with h_1 nor with h_2 then it is clear that $v'(t) \leq 0$. If t is comparable only with one of h_1, h_2 , then it is comparable either with h^- or with h^+ , and also $v'(t) \leq 0$. Let t be comparable with h_1 and with h_2 . Suppose $t < h_1$ (a case $t > h_1$ is analogous) then $t < h_2$. ($h_1 < t < h_2$ is impossible, so $t < h^+$). Then if t is comparable also with h^- then $v'(t) = 0$, and in the opposite case $S \supset W^4 = \{t, h_2, h^-, h_1\}$. ■

Lemma 10. *If $S \supseteq W^{2k}$ ($k \geq 2$) and form f_S is positive semidefinite then $C(S) \neq \emptyset$.*

Let $t \in T = S \setminus W^{2k}$. Put $S^-(t) = |\{s_i^- | t < s_i^-\}| \cup \{s_i^- | t > s_i^-\}|$, $S^+(t) = |\{s_i^+ | t < s_i^+\}| \cup \{s_i^+ | t > s_i^+\}|$. We prove that positive semidefiniteness of f_S implies $S^-(t) = S^+(t)$.

Really, let $S^-(t_0) > S^+(t_0)$ for fixed $t_0 \in T$ (a case $S^-(t_0) < S^+(t_0)$ is analogous). Consider $x : S \rightarrow \mathbb{R}_n$ where $x(s_i^-) = -1$, $x(s_i^+) = 1$, ($i = \overline{1, k}$), $x(t_0) = \varepsilon$, $0 < \varepsilon < 1$ and $x(t) = 0$ for $t \in T \setminus \{t_0\}$. It is easy to see that $f_S(x) < 0$.

Now consider vector $v : S \rightarrow \mathbb{R}_n$ where $v(s_i^-) = -1$, $v(s_i^+) = 1$, $v(t) = 0$ for $t \in T$. $S^-(t) = S^+(t)$ ($t \in T$) implies $v'(s) = 0$ for any $s \in S$. It is clear, that $v \in H_n$ and therefore $v \in C(S)$. ■

Proposition 6. *If S is an antimonotonous poset and form f_S is positive semidefinite then $\Gamma(S) = A_n$.*

If S is cyclical, then Lemmas 8,9,10 imply the statement. If S is acyclical, then by Proposition 4 the Tits form $\mathcal{T}_S$ is positive semidefinite, so $\Gamma(S)$ is one of $A_n, D_n, E_6, E_7, E_8, \tilde{D}_n, \tilde{E}_6, \tilde{E}_7, \tilde{E}_8$ ($\Gamma(\tilde{A}_n)$ is cyclical). If $\Gamma(S) \neq A_n$ then Proposition 5, Examples 2 and 5 and Lemma 7 imply the existence of the Dynkin point and, by Lemma 3 (and Remark 2), $C(S) \neq \emptyset$.

6. We consider now $\Gamma(S) = A_n$. In this case poset S up to antiisomorphism is defined by its order and by subset $S^\times$ consisting of junction points (see s .3). It is clear that $S^\times = \emptyset$ if and only if S is a chain.

Let poset $W^{k,k+1} = \{s_1^-, \dots, s_k^-, s_1^+, \dots, s_{k+1}^+ | s_i^- < s_i^+, s_i^- < s_{i+1}^+, i = \overline{1, k}\}$, $W^{k+1,k} = \{s_1^-, \dots, s_{k+1}^-, s_1^+, \dots, s_k^+ | s_i^+ > s_i^-, s_i^+ > s_{i+1}^-, i = \overline{1, k}\}$.

Lemma 11. *If $\Gamma(S) = A_n$ and $\Gamma(S) \supseteq W$ of the form $W^{k,k+1}$ (resp. $W^{k+1,k}$), moreover $s_1^+, s_{k+1}^+ \notin S^\times$ (resp. $s_1^-, s_{k+1}^- \notin S^\times$) then $C(S) \neq \emptyset$.*

Let, for the sake of definiteness, $S \supset W^{k,k+1}$. Consider vector v such that $v(s_i^-) = -2$, $i = \overline{1, k}$; $v(s_i^+) = +2$, $i = \overline{2, k}$; $v(s_1^+) = v(s_{k+1}^+) = 1$; $v(t) = 0$ for $t \in S \setminus W^{k,k+1}$. We prove that $v \in C(f)$.

Indeed, $v \in H_n$; $v'(s_1^-) = v'(s_k^-) = -1$; $v'(s_i^-) = 0$ for $i = \overline{2, k-1}$; $v'(s_i^+) = 0$ for $i = \overline{1, k+1}$. Absence of branch points implies that if $t \notin W$ is comparable with $w \in W$ then $w \in \{s_1^+, s_{k+1}^+\}$. If t is comparable with both s_1^+, s_{k+1}^+ then S would be cyclical. $s_1^+, s_{k+1}^+ \notin S^\times$ implies that it can be only $t > w$. Therefore every t either is comparable

exactly with one s_i^- and one s_i^+ , or is not comparable with any $w \in W$, hereof $v'(t) \leq 0$. ■

We call a poset ζ a *wattle* [3] if it is the union of not intersecting chains Z_i , $|Z_i| \geq 2$, $i = \overline{1, t}$, $t > 1$, in which the minimal element of Z_i is less than the maximal element of Z_{i+1} and there are no other comparisons between elements of different Z_i . $\Gamma(\zeta) = A_n$. According to [3] we denote $\zeta = \langle n_1, \dots, n_t \rangle$ where $n_i = |Z_i|$.

For a poset S we consider a disconnected subgraph $\Gamma(S^\times)$ of $\Gamma(S)$. Denote $S_i^\times$ its connected components.

Lemma 12. *A poset S where $\Gamma(S) = A_n$ is either a chain or a wattle if (and only if) the orders of all $S_i^\times$ are even.*

If S is a wattle then the statement is evident (and we will not use it). We will prove the converse statement induction by $|S|$. The base is evident. Let $|S| = n + 1$. $\Gamma(S) = \dots s_{n-1} \text{---} s_n \text{---} s_{n+1}$ where s_{n+1} is a terminal point (therefore $s_{n+1} \notin S^\times$). For the sake of definiteness we will suppose that $s_n > s_{n+1}$. So s_{n+1} is minimal. Put $S' = S \setminus \{s_{n+1}\}$ and $S'' = S \setminus \{s_{n+1}, s_n\}$. We have two possibilities: 1) $s_{n-1} > s_n$; 2) $s_{n-1} < s_n$.

1) $s_n \notin S^\times$, $(S')^\times = S^\times$. By assumption of the induction S' is a wattle in which s_n is a minimal terminal point. It is clear that S is either a wattle or a chain. (If $S' = \langle n_1, \dots, n_t \rangle$ then $S' = \langle n_1, \dots, n_t + 1 \rangle$).


2) $s_n \in S^\times$ (S' does not satisfy the induction's assumption!). $s_n \in S_p^\times$, $|S_p^\times| \equiv 0 \pmod{2}$. So $s_{n-1} \in S_p^\times \subset S^\times$. s_{n-1} is a terminal point of S'' and so $s_{n-1} \notin (S'')^\times$. If $S^\times = \bigcup_{i=1}^p S_i^\times$ then $(S'')^\times = \bigcup_{i=1}^{p-1} S_i^\times \cup (S_p^\times \setminus \{s_n, s_{n+1}\})$.

Consequently S'' satisfies the induction's assumption and so it is either a chain or a wattle, in which s_{n-1} is a minimal point. If $S'' = \langle n_1, \dots, n_t \rangle$ then $S = \langle n_1, \dots, n_t, 2 \rangle$. ■

Proposition 7. *If a form f_S is positive semidefinite and $C(S) = \emptyset$, then S is either a chain or a wattle.*

Proposition 6 implies $\Gamma(S) = A_n$.

If S is neither a chain nor a wattle then by Lemma 12 there exists $S_p^\times$ such that $|S_p^\times| \equiv 1 \pmod{2}$. It is easy to see that $S_p^\times$ is $W^{k,k+1}$ or $W^{k+1,k}$. Let, for the sake of definiteness $S_p^\times = W^{k,k+1} =$




and $S_p^\times$ is a connected component in $S^\times$, imply that there exists $s_0^- \in S \setminus S^\times$ such that $s_0^- \rightarrow s_1^+$. Analogously there exists $s_{k+1}^- \in S \setminus S^\times$, $s_{k+1}^- \rightarrow s_{k+1}^+$. It is easy to see that $S_p^\times \cup \{s_0^-, s_{k+1}^-\} = W^{k+2,k+1}$ and $C(S) \neq \emptyset$ by Lemma 11. ■

Example 4 (s. 3) implies that the condition of positive semidefiniteness of f_S can not, generally speaking, be excluded.

Hypothesis. *If S is acyclical and $\Gamma(S) \neq A_n$ then $C(S) \neq \emptyset$.*

In some cases the existence of $v \in C(f)$ for acyclical S is evident. We give however the example of an acyclic poset S and $v \in C(S)$ which we constructed only by means of computer.

Example 6.



7. Let $\zeta = \langle n_1, \dots, n_t \rangle$ ($t > 1$) be a wattle, where $n_i = |Z_i|$, $\sum_{i=1}^t n_i = n$, $n_i > 1$, $i = \overline{1, t}$.

In [3] the minimal points of chains Z_i , $i = \overline{1, t-1}$ are denoted by z_i^- , and maximal points of chains Z_i , $i = \overline{2, t}$ are denoted by z_i^+ , $z_i^- < z_{i+1}^+$. The rest (i. e. not junction) points are called *common* (including the maximal point of chain Z_1 and a minimal point of chain Z_t). They are compared only with points from its chain.

A *width* $\omega(S)$ of a partially ordered set S is the maximal number of its pairwise uncomparable elements. We attach to any poset S a rational number $r(S) = \frac{n+1}{t} - 1$ where $n = |S|$, $t = \omega(S)$. If S is a chain then $w(s) = 1$, $r(S) = n$.

It is clear that there exist many wattles with the same r . We prove below, however, that to any noninteger $r > 1$ it is corresponded exactly one (uniform in sense [3]) P -faithful (= antimonotonous, see Corollary 3) wattle, which we will call here r -wattle and denote $\zeta(r)$.

For a positive rational a we put $\{a\} = a - [a]$. Let r be a positive noninteger rational number greater than 1. q/t is the representation of $\{r\}$ in the form of irreducible fraction. We write out the sequence of integers $n_1, \dots, n_t$, which will be the orders of sets Z_i in $\zeta(r)$. Put $n_1 = n_t = [r] + 1$, $n_i = [ri] - [r(i-1)] + 1$ for $i = \overline{2, t-1}$. It is clear, that $[\{r\}i] - [\{r\}(i-1)]$ is either 1 or 0. Therefore n_i is either $1 + [r]$ or $2 + [r]$. The number of those i for which $n_i = 2 + [r]$ is $q - 1$; $n = t([r] + 1) + q - 1$, $r(\zeta(r)) = r$.

Remark, that r -wattles are uniform in sense of [3] and conversely.

Thus, we corresponded to each noninteger rational number $r > 1$ a wattle $\zeta(r)$. We can consider also integer numbers, putting for natural r that $\zeta(r)$ is a chain of length r . We will call r -sets all posets of form $\zeta(r)$ $r \geq 1$ (i. e. uniform wattles and chains).

Theorem. Let form f_S be positive semidefinite ($\Gamma(S)$ be connected). Then $C(S) = \emptyset$, if and only if S is r -set.

If r is integer then the statement is evident (see [3]). Therefore as a matter of fact we need to prove, counting Proposition 7, that $C(\zeta) = \emptyset$ if and only if ζ is r -wattle.

We attach to r -wattle $\zeta(r)$ a vector $x : \zeta \rightarrow \mathbb{R}^+$, $x(s) = 1$ for $s \in \zeta \setminus \zeta^\times$, $x(z_i^-) = \{ir\}$, $x(z_i^+) = 1 - x(z_{i-1}^-)$. $(q, t) = 1$ implies $x(s) > 0$ for any $s \in \zeta(r)$.

It is possible to check that $x \in \text{St}(\zeta)$ either spontaneously, using the definition of $\text{St}(\zeta)$, or using the following lemma.

Lemma 13 (see [3], Lemma 5). *Vector $x : \zeta \rightarrow \mathbb{R}$ is contained in $\text{St}(\zeta)$ if and only if there exist such positive α, β that*

- 1) $x(s) = \alpha$ for $s \in \zeta \setminus \zeta^\times$ (we can suppose $\alpha = 1$ multiplying x by $\lambda \in \mathbb{R}^+$);
- 2) $x(z_i^-) + x(z_{i+1}^+) = \alpha$ for $i = \overline{1, t-1}$;
- 3) $\sum_{s \in Z_i} x(s) = \beta, i = \overline{1, t}$.

Proof of the lemma is almost evident. Remark merely that at first 2) should be proved (it follows from $\frac{\partial f_s}{\partial z_i^-}(x) = \frac{\partial f_s}{\partial z_i^+}(x), i = \overline{2, t-1}$, and then 1).

Vector x constructed above evidently satisfies the conditions 1), 2) of Lemma 13. It is easy to check (for $\alpha = 1$) that

$$\sum_{s \in Z_i} x(s) = r(i = \overline{1, t}); \quad (4)$$

$$x'(s) = 1 + r; \quad (5)$$

$$\sum_{s \in \zeta(r)} x(s) = tr, \quad (6)$$

Thus, $x \in \text{St}(\zeta)$, so $\tilde{\text{St}}(\zeta) \neq \emptyset$, and $C(S) = \emptyset$ by Corollary 3.

It remains to prove that any P -faithful wattle ζ is r -wattle (where $[z] = |Z_1| - 1, \{r\} = x(z_1^-)$). This follows from the next statement.

Lemma 14. *Let $\zeta = \langle z_1, \dots, z_t \rangle$ and $\hat{\zeta} = \langle \hat{z}_1, \dots, \hat{z}_t \rangle$ are two P -faithful wattles, $x \in \text{St}(\zeta), \hat{x} \in \text{St}(\hat{\zeta})$ ($\hat{\alpha} = \alpha = 1$). Then, if $Z_1 = \hat{Z}_1$ and $x(s) = \hat{x}(s)$ for $s \in Z_1 = \hat{Z}_1$ then $\zeta = \hat{\zeta}$ and $x(s) = \hat{x}(s)$ for $s \in \zeta$.*

It is sufficient to prove that if $m \leq \max\{t, \hat{t}\}$, then $z_i = \hat{z}_i$ for $i \leq m$ and $x(s) = \hat{x}(s)$ for $s \in \cup_{i=1}^m Z_i$. Lemma 13 implies this by induction with respect to m (see [3]). ■ ■

There was introduced the numerical function $\rho(r) = 1 + \frac{r-1}{r+1}$ where $r \in \mathbb{N}$ [3]. We spread this definition on the case $r \geq 1$ is rational. Put $\rho(r_1, \dots, r_t) = \sum_{i=1}^t \rho(r_i)$. If Z_n is a chain of

order n then $P(Z_n) = \rho(n)$ [3]. Let $\zeta(r)$ be a wattle. Vector $\bar{x} = (tr)^{-1}x \stackrel{(6)}{\in} P_n \cap \text{St}(\zeta(r))$ (where x is a vector constructed in the proof of the theorem).

$$\begin{aligned} P(\zeta(r)) &= f_{\zeta(r)}^{-1}(\bar{x}) = (tr)^2 f_{\zeta(r)}^{-1}(x) \stackrel{(3)}{=} \\ &= 2(tr)^2 \left(\sum_{s \in \zeta(r)} x'(s)x(s) \right)^{-1} \stackrel{(5), (6)}{=} \frac{2t^2 r^2}{(1+r)tr} = \frac{2tr}{1+r} = t\rho(r). \end{aligned}$$

This formula is true if $t = 1$ (i. e. in the case of chain). For any positive rational $r = \frac{l}{t}$ ($(l, t) = 1$) $t\rho(r) = \frac{2lt}{l+t}$. Introduce the function $P(r) = \frac{2lt}{l+t}$ (for $n \in \mathbb{N}$ $P(n) = \rho(n)$). Thus, for any $r \geq 1$

$$P(\zeta(r)) = t\rho(r) = P(r). \quad (7)$$

Appendix. We call a poset S *connected* if graph $\Gamma(S)$ is connected. The theorem and Corollary 3 imply that a connected poset S is P -faithful if and only if it is r -set [3–7]. On the other hand, [3–7] imply our theorem only if f_S is positive definite (but not positive semidefinite).

We remind about role of P -faithful posets in representation theory. We will write $S = S_1 \sqcup S_2$, if $S = S_1 \cup S_2$, $S_1 \cap S_2 = \emptyset$ and elements S_1 are not comparable with elements S_2 . $S = Z_1 \sqcup \dots \sqcup Z_p$ is *primitive*, if Z_i are chains, $i = \overline{1, p}$. We denote it $(n_1, \dots, n_p)$ if $n_i = |Z_i|$. The characterization of antimonotonous disconnected posets follows from the theorem and Lemma 2.

Any poset $S = \bigsqcup_{i=1}^p S_i$ where S_i are connected components. According to Lemma 5, $P(S) = \sum_{i=1}^p P(S_i)$, and if S is primitive then $P(S) = \sum_{i=1}^p \rho(n_i) = \rho(n_1, \dots, n_t)$.

A role of quadratic forms in the theory of representations of quivers and posets is well-known [12].

The norm of a relation $\|S, \leq\| = \inf_{u \in \overline{P}_n} f_S(u)$ was introduced in [2] on base of form f_S . In view of Lemma 5 it is naturally to consider instead of $\|S, \leq\|$ the function $P(S) = \|S, \leq\|^{-1}$ [2].

Proposition 8. S has finite (respectively tame) type if and only if $P(S) < 4$ (respectively $P(S) = 4$).

With this point of view the Kleiner's list of the critical posets [8] is the list of P -faithful posets S_i , for which $P(S) = 4$.

4 posets of Kleiner's list are primitive:

$$(I). \quad (1, 1, 1, 1), \quad (2, 2, 2), \quad (1, 3, 3), \quad (1, 2, 5), \quad \text{and the fifth is} \quad (4) \sqcup K,$$

where $K = \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \begin{array}{c} \bullet \\ / \backslash \\ \bullet \end{array} = \langle 2; 2 \rangle = \zeta(1\frac{1}{2})$. It is easy to see that any chain is P -faithful, and s. 7 implies that K is also P -faithful ($P(K) = 2, 4$). By Lemma 5 a disconnected poset is P -faithful if and only if all its components are P -faithful.

The list of critical sets [10]:

$$(II). \quad (1, 1, 1, 1, 1), \quad (1, 1, 1, 2), \quad (2, 2, 3), \quad (1, 3, 4), \quad (1, 2, 6), \quad (6) \sqcup K$$

can be characterized as the list of S , having the following properties:

- 1) $P(S) > 4$,
- 2) if $S' \subset S$ then $P(S') \leq 4$.

The following statement play central role in the theory of representations of posets [8], [10].

A poset S is finitely represented (respectively tame), if and only if S does not contain subsets I (respectively II).

It has been naturally to suppose that all P -faithful poset are either chains or some sets, for which K is the least representative. This was a reason to introduce P -faithful posets [3].

Now we show how the lists (I), (II) can be obtained using characteristic of (connected) P -faithful sets and formula (7). It is easy to check that $P(S) = 4$ for $S \in \text{I}$ and $P(S) > 4$ for $S \in \text{II}$ (in view of Lemma 5 and formula (7)).

We call a P -faithful poset S *utmost* if $P(S) \geq 4$ and $P(S') \leq 4$ for any $S' \subset S$ (hereat S' can be supposed P -faithful).

Lemma 15. *Not primitive utmost S is of the form $K \sqcup Z_m$ where m is 4 or 5.*

Let S contains a connected component $\zeta(r)$ where $\{r\} = \frac{q}{t}$, $t > 1$, $q < t$, $(q, t) = 1$. The characterization of P -faithful posets implies $\omega(S) < 4$ since in the opposite case $S \supset S' = (2, 1, 1, 1)$, $\rho(2, 1, 1, 1) = 4\frac{1}{3} > 4$. Consequently $t \leq 3$, moreover if $t = 3$ then $\zeta(r) = S$.

Let $t = 3$, $1 \leq q \leq 2$. If $[r] \geq 2$ then $S \supset S' = S \setminus \{z_1^-, z_3^+\}$. S' is a primitive poset containing $(2, 3, 2)$, $\rho(2, 2, 3) = 4\frac{1}{6}$, $\rho(S') > 4$. If $[r] = 1$ then either $r = 1\frac{1}{3}$ or $r = 1\frac{2}{3}$. We obtain $\rho(r) \leq 1\frac{1}{4}$, then $P(S) = 3\rho(r) < 4$ (see (7)).

Let $t = 2$ and $S \neq \zeta(1\frac{1}{2}) \sqcup \hat{S}$. If $S = \zeta(r)$ then $P(S) = 2\rho(r) < 4$ since $\rho(r) < 2$ for any r . So, $S = \zeta(r) \sqcup \hat{S}$, $r > 1\frac{1}{2}$, i. e. $r \geq 2\frac{1}{2}$, $\zeta(r) \supset \{\zeta(r) \setminus z_1^-\} \supseteq (2, 3)$. For $|\hat{S}| > 1$, $S \supset S'$, contains $(2, 2, 3)$ or $(1, 1, 2, 3)$ and $P(S') > 4$. Hence $|\hat{S}| = 1$. Then $[r] < 3$, since otherwise $\zeta(r) \supset S' = \zeta(r) \setminus z_1^- \supseteq (3, 4)$ and $P(S' \sqcup (1)) > 4$ since $\rho(3, 4, 1) > 4$. For $[r] = 2$ we obtain $P(S) < 4$ since $P(\zeta(2\frac{1}{2})) = 2 \cdot 1\frac{3}{7}$, $P(\zeta(2\frac{1}{2}) \sqcup (1)) = 2\frac{6}{7} + 1$ (Lemma 5).

Let, finally, $S = \zeta(1\frac{1}{2}) + \hat{S}$ ($S \neq \zeta(1\frac{1}{2})$) since $P(\zeta(1\frac{1}{2})) = 2, 4$, then if $w(\hat{S}) > 1$ then $S \supset S' = \{\zeta(1, \frac{1}{2}) \setminus z_1^-\} \sqcup (1, 1) = (2, 1, 1, 1)$, $\rho(S') > 4$. If $\hat{S} = Z_m$ then for $m < 4$ $\rho(m) < 1, 6$ and $P(S) < 4$, and for $m > 5$, $S \supset S' = (\zeta(1\frac{1}{2}) \sqcup Z_5)$, $\rho(S') > 4$, $P(K \sqcup Z_4) = 4$, $P(K \sqcup Z_5) = 4\frac{1}{15}$. ■

Proposition 9. *P -faithful S is utmost if and only if $S \in \text{I} \cup \text{II}$.*

If S is not primitive then Lemma 15 implies the statement. Let S is primitive. Then $w(S) > 2$ and $S \notin \{(1, 1, n), (1, 2, 2), (1, 2, 3), (1, 2, 4)\}$ (otherwise $P(n_1, \dots, n_t) = \rho(n_1, \dots, n_t) < 4$). In the rest cases we can see that if $S \notin \text{I} \cup \text{II}$ then $S \supset S' \in \text{II}$, and if $S \in \text{I} \cup \text{II}$ then $S \not\supset S' \in \text{II}$. ■

Propositions 8, 9 (in view of $P(S) = 4$ for $S \in \text{I}$ and $P(S) > 4$ for $S \in \text{II}$) imply the main theorems [8] and [10].

P -faithful posets, for which $P = 4$, play important role in representation theory. We don't know whether P -faithful posets with $P = n > 4$ play some analogous role. In [14] primitive posets with $P = 5$ are written out. We give the example (probably unique) of a not primitive poset $S = \zeta(3\frac{1}{2}) \sqcup (17)$ for which $P(S) = 5$ (see (7) and Lemma 3).

It is Example 4 (s. 3) where $C(S) = \emptyset$ but S is not P -faithful. We hope that studying of $C(S)$ can be interesting for the representation theory. Remark, that in [2] the norm $\|P\|$ of an arbitrary binary relation P (on finite set) and corresponding notion of P -faithfulness are defined (these notions can be used for locally scalar representations (see [9]) in Hilbert spaces). However in this case it is more complicate to review P -faithful sets. Such investigation would be seemingly rather difficult and interesting problem.

References

- [1] Fihthengolts G.M. Kurs differentsialnogo i integralnogo ischisleniya, V. 1. — Moscow–Leningrad, 1947. — P. 690.
- [2] Roiter A.V. The norm of a relation / Representation Theory. I. Finite Dimensional Algebras, Proc., Ottawa, 1984 / Lecture Notes in Math., 1177. — P. 269–272.
- [3] Nazarova L.A., Roiter A.V. Norma otnosheniya, razdelyayushchie funktsii i predstavleniya markirovannykh kolchanov // Ukr. Math. Jour. — 2002. — **54**, N 6. — P. 808–840.
- [4] Zeldich M.V. Pro harakteristiki formy chastkovo vporyadkovanykh mnozhyn z odnozvyaznym grafom Hasse. // Visnyk Kyivskogo universytetu (serie: phys.-math.sci.). — 2001. — N 4. — P. 36–44.
- [5] Zeldich M.V. Pro ρ -tochni chastkovo vporyadkovani mnozhyny // Visnyk Kyivskogo universytetu (serie: phys.-math.sci.). — 2001. — N 4. — P. 45–51.
- [6] Sapelkin A.I. P -tochnye chastichno uporyadochennyye mnozhestva // Ukr. Math. Jour. — 2002. — **54**, N 10. — P. 1381–1396.
- [7] Zeldich M.V. O harakteristicheskikh i kratno tranzitivnykh formakh chastichno uporyadochennyykh mnozhestv. O P -tochnykh chastichno uporyadochennykh mnozhestvakh / Preprint, Kievskiy natsionalnyy universitet im. Tarasa shevchenka, Kiev, 2002, 64 p.
- [8] Kleiner M.M. Chastichno uporyadochennyye mnozhestva konechnogo tipa // Zap. nauchn. seminarov LOMI AN SSSR. — 1972. — **28**. — P. 32–42.
- [9] Ringel C.M. Tame Algebras and Integral Quadratic Forms, Springer-Verlag 1099, 1984, P. 376.
- [10] Nazarova L.A. Chastichno uporyadochennyye mnozhestva beskonechnogo tipa // Izv. AN SSSR, sektsiya matem. — 1975. — **39**, N 5. — P. 963–991.
- [11] Zavadskiy A.G., Nazarova L.A. Chastichno uporyadochennyye mnozhestva ruchnogo tipa / In book "Matrichnye zadachi". — Kiev, In-t of Mathematics AN USSR, 1977. — P. 122–143.
- [12] Gabriel P., Roiter A.V. Representations of finite-dimensional algebras. — Springer-Verlag, Algebra VIII, 1992.
- [13] Kruglyak S.A., Roiter A.V. Lokalno-skalyarnie predstavleniya grafov v kategorii gilbertovykh prostranstv // Func. Analis i Pril. — 2005. — N 2.
- [14] Redchuk I.K., Roiter A.V. Singulyarnyye lokalno-skalyarnyye predstavleniya v gilbertovykh prostranstvakh i razdelyayushchie funktsii // Ukr. Math. Jour. — 2004. — **56**, N 6. — P. 796–809.